

1.3.1	Entecavir
SPC, Labeling and Package Leaflet	NL-Netherlands

Package leaflet: Information for the user

Entecavir Krka 0,5 mg filmomhulde tabletten Entecavir Krka 1 mg filmomhulde tabletten entecavir

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Entecavir Krka is and what it is used for
2. What you need to know before you take Entecavir Krka
3. How to take Entecavir Krka
4. Possible side effects
5. How to store Entecavir Krka
6. Contents of the pack and other information

1. What Entecavir Krka is and what it is used for

Entecavir Krka tablets are anti-viral medicines, used to treat chronic (long term) hepatitis B virus (HBV) infection in adults. Entecavir Krka can be used in people whose liver is damaged but still functions properly (compensated liver disease) and in people whose liver is damaged and does not function properly (decompensated liver disease).

Entecavir Krka tablets are also used to treat chronic (long term) HBV infection in children and adolescents aged 2 years to less than 18 years. Entecavir Krka can be used in children whose liver is damaged but still functions properly (compensated liver disease).

Infection by the hepatitis B virus can lead to damage to the liver. Entecavir Krka reduces the amount of virus in your body, and improves the condition of the liver.

2. What you need to know before you take Entecavir Krka

Do not take Entecavir Krka:

- **if you are allergic (hypersensitive)** to entecavir or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Entecavir Krka

- **if you have ever had problems with your kidneys**, tell your doctor. This is important because Entecavir Krka is eliminated from your body through the kidneys and your dose or dosing schedule may need to be adjusted.
- **do not stop taking Entecavir Krka without your doctor's advice** since your hepatitis may worsen after stopping treatment. When your treatment with Entecavir Krka is stopped, your doctor

1.3.1	Entecavir
SPC, Labeling and Package Leaflet	NL-Netherlands

will continue to monitor you and take blood tests for several months.

- **discuss with your doctor whether your liver functions properly** and, if not, what the possible effects on your Entecavir Krka treatment may be.
- **if you are also infected with HIV** (human immunodeficiency virus) be sure to tell your doctor. You should not take Entecavir Krka to treat your hepatitis B infection unless you are taking medicines for HIV at the same time, as the effectiveness of future HIV treatment may be reduced. Entecavir Krka will not control your HIV infection.
- **taking Entecavir Krka will not stop you from infecting other people with hepatitis B virus (HBV)** through sexual contact or body fluids (including blood contamination). So, it is important to take appropriate precautions to prevent others from becoming infected with HBV. A vaccine is available to protect those at risk from becoming infected with HBV.
- **Entecavir Krka belongs to a class of medicines that can cause lactic acidosis** (excess of lactic acid in your blood) and enlargement of the liver. Symptoms such as nausea, vomiting and stomach pain might indicate the development of lactic acidosis. This rare but serious side effect has occasionally been fatal. Lactic acidosis occurs more often in women, particularly if they are very overweight. Your doctor will monitor you regularly while you are receiving Entecavir Krka.
- **if you have previously received treatment for chronic hepatitis B**, please inform your doctor.

Children and adolescents

Entecavir Krka should not be used for children below 2 years of age or weighing less than 10 kg.

Other medicines and Entecavir Krka

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Entecavir Krka with food and drink

In most cases you may take Entecavir Krka with or without food. However, if you have had a previous treatment with a medicine containing the active substance lamivudine you should consider the following. If you were switched over to Entecavir Krka because the treatment with lamivudine was not successful, you should take Entecavir Krka on an empty stomach once daily. If your liver disease is very advanced, your doctor will also instruct you to take Entecavir Krka on an empty stomach. Empty stomach means at least 2 hours after a meal and at least 2 hours before your next meal.

Children and adolescents (from 2 to less than 18 years of age) can take Entecavir Krka with or without food.

Pregnancy, breast-feeding and fertility

Tell your doctor if you are pregnant or planning to become pregnant. It has not been demonstrated that Entecavir Krka is safe to use during pregnancy. Entecavir Krka must not be used during pregnancy unless specifically directed by your doctor. It is important that women of childbearing age receiving treatment with Entecavir Krka use an effective method of contraception to avoid becoming pregnant.

You should not breast-feed during treatment with Entecavir Krka. Tell your doctor if you are breast-feeding. It is not known whether entecavir, the active ingredient in Entecavir Krka, is excreted in human breast milk.

Driving and using machines

Dizziness, tiredness (fatigue) and sleepiness (somnolence) are common side effects which may impair your ability to drive and use machines. If you have any concerns consult your doctor.

1.3.1	Entecavir
SPC, Labeling and Package Leaflet	NL-Netherlands

Entecavir Krka contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Entecavir Krka

Not all patients need to take the same dose of Entecavir Krka.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

For adults the recommended dose is either 0.5 mg or 1 mg once daily orally (by mouth).

Your dose will depend on:

- whether you have been treated for HBV infection before, and what medicine you received.
- whether you have kidney problems. Your doctor may prescribe a lower dose for you or instruct you to take it less often than once a day.
- the condition of your liver.

For children and adolescents (from 2 to less than 18 years of age), your child's doctor will decide the right dose based on your child's weight. An entecavir oral solution is recommended for patients weighing from 10 kg to 32.5 kg.

Children weighing at least 32.6 kg may take an entecavir oral solution or the 0.5 mg tablet. All dosing will be taken once daily orally (by mouth). There are no recommendations for entecavir in children less than 2 years of age or weighing less than 10 kg.

For children and adolescents (from 2 to less than 18 years of age), <Invented name> 0.5 mg tablets are available.

Your doctor will advise you on the dose that is right for you. Always take the dose recommended by your doctor to ensure that your medicine is fully effective and to reduce the development of resistance to treatment. Take Entecavir Krka as long as your doctor has told you. Your doctor will tell you if and when you should stop the treatment.

Some patients must take Entecavir Krka on an empty stomach (see **Entecavir Krka with food and drink** in **Section 2**). If your doctor instructs you to take Entecavir Krka on an empty stomach, empty stomach means at least 2 hours after a meal and at least 2 hours before your next meal.

If you take more Entecavir Krka than you should

Contact your doctor at once.

If you forget to take Entecavir Krka

It is important that you do not miss any doses. If you miss a dose of Entecavir Krka, take it as soon as possible, and then take your next scheduled dose at its regular time. If it is almost time for your next dose, do not take the missed dose. Wait and take the next dose at the regular time. Do not take a double dose to make up for a forgotten dose.

Do not stop Entecavir Krka without your doctor's advice

Some people get very serious hepatitis symptoms when they stop taking Entecavir Krka. Tell your doctor immediately about any changes in symptoms that you notice after stopping treatment.

1.3.1	Entecavir
SPC, Labeling and Package Leaflet	NL-Netherlands

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Patients treated with entecavir have reported the following side effects:

Adults

- common (at least 1 in 100 people): headache, insomnia (inability to sleep), fatigue (extreme tiredness), dizziness, somnolence (sleepiness), vomiting, diarrhoea, nausea, dyspepsia (indigestion), and increased blood levels of liver enzymes.
- uncommon (at least 1 in 1,000 people): rash, hair loss.
- rare (at least 1 in 10,000 people): severe allergic reaction.

Children and adolescents

The side effects experienced in children and adolescents are similar to those experienced in adults as described above with the following difference:

Very common (at least 1 in 10 patients): low levels of neutrophils (one type of white blood cells, which are important in fighting infection).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Entecavir Krka

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the packaging after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Entecavir Krka contains

- The active substance is entecavir.
Each film-coated tablet contains 0.5 mg entecavir (as monohydrate).
Each film-coated tablet contains 1 mg entecavir (as monohydrate).
- The other ingredients are lactose monohydrate, microcrystalline cellulose (E 460), crospovidone type B (E 1202), hypromellose (E 464) and magnesium stearate (E 470b) in the tablet core and hypromellose (E 464), titanium dioxide (E 171), macrogol 400 (E 1521), polysorbate 80 – only for 0.5 mg film-coated tablets and iron oxide red (E 172) – only for 1 mg film-coated tables in film coating.

1.3.1	Entecavir
SPC, Labeling and Package Leaflet	NL-Netherlands

See section 2 "Entecavir Krka contains lactose".

What Entecavir Krka looks like and contents of the pack

White triangular shaped film coated tablets debossed 'A' on one side and '88' on the other side of 8.7 x 8.4 mm.

Pink triangular shaped film coated tablets debossed 'A' on one side and '89' on the other side of 10.9 x 10.5 mm.

Entecavir Krka 0.5 mg film-coated tablets are supplied in boxes containing 30 or 90 film-coated tablets in OPA/Aluminium/PVC-Aluminium blisters.

Entecavir Krka 1 mg film-coated tablets are supplied in boxes containing 30 or 90 film-coated tablets in OPA/Aluminium/PVC-Aluminium blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenië

Manufacturers

Pharmacare Premium Ltd., HHF003 Hal Far Industrial Estate, Hal Far, BBG 3000 Birzebbugia, Malta

This drug is registered in the registry under:

RVG 119614 Entecavir Krka 0,5 mg filmomhulde tabletten

RVG 119615 Entecavir Krka 1 mg filmomhulde tabletten

This medicinal product is authorised in the Member States of the European Economic Area under the following names:

Naam van de lidstaat	Naam van het geneesmiddel
Nederland, België, Roemenië	Entecavir Krka
Duitsland	Entecavir TAD
Frankrijk, Oostenrijk	Entecavir HCS
Bulgarije	Ентекавир Крка
Slovenië	Entekavir Krka

Deze bijsluiter is voor het laatst goedgekeurd in november 2021.